

Detection of starquakes on magnetically active red dwarfs

M.E. Contadakis¹, S.J. Avgoloupis², J. H. Seiradakis^{3,+}, Ch. Papantoniou¹

¹Department of of Surveying and Geodesy, Aristotle University of Thessaloniki, GR-54124

²Faculty of Education, Schoole of Primary Education, Aristothe University of Thessaloniki

³Department of Physics, Aristotle University of Thessaloniki, GR-54124, Thessaloniki Greece.

Abstract: The results of the analysis of one colour (B, or U) observations of the Stephanion Observatory of different red dwarfs (EV Lac, BY Drac, AD Leo, YZ CMin) at the quiescence stage of their activity, indicate that transient high frequency oscillations occur. The observed oscillations are of the same frequencies range of the sunquakes.

1 Introduction

The scientific team of the Stephanion Observatory, University of Thessaloniki contributed to the research of high frequency optical oscillations on red dwarfs by participating in international programs for Multiwavelength observation of strong Flares of selected flare stars ([12]). These joined research shed plenty of light on the phenomenon of high frequency optical oscillations. Nevertheless a better understanding of the high-frequency oscillations demand a unified analysis of the flare light-curve for a wider time window covering pre-flare, flare and post flare and a broader band of frequencies. Thus in addition to the international campaign research the Stephanion Observatory group observe and analysis one colour (B, or U) observations of the Stephanion Observatory of different red dwarfs: EV Lac([1], [2] and [7]), AD Leo ([4] and [5]),YZ CMin ([3],[9]), V 390 Auri ([6],[10]), UV Cet([8]), at any stage of their activity (quiescence, weak flares, strong flares). In this paper we present the analysis of the quiet state observations of the stars EV Lac, BY Drac , AD Leo, YZ Cmin in order to realize if starquakes appear far apart from the observed flares, during the quiet state of the stars, as a result of the general magnetic activity of the star.

2 The Observations

The observations were carried out with the help of the 30-inch Cassegrain telescope of the Stephanion Observatory ($\phi = 22^{\circ}49'45''$, $\lambda = 37^{\circ}45'09''$ and $h = 900m$) which is equipped with a Johnson photometer. The digitized recording system has a recording resolution of 0.108s. This facility allows the reliable analysis of the observational data for a wide range of frequencies (4.63Hz to 4.63×10^{-3} Hz i.e. periods 0.216s to 216s) and permits the reliable spectral analysis of small part of the light-curve. The observations were made mostly in B and occasionally in U color during the years 2002 and 2004 . Selected parts of the quiescence state were analyzed in order to detect high frequency optical oscillations.

3 Analysis

The data consist of a sample of relative intensities i.e. $(I - I_o)/I_o$, where I is the flare intensity and I_o is the quiet star intensity. The extent, as well as the resolution, of the observational sample permit a reliable Discrete Fourier Transform analysis of sufficient light-curve parts for the determination of possible transient oscillations in the frequency domain and their approximate location in the time domain. So the analysis comprises the following steps: (1) With the help of DFT-analysis we deduce

the power spectra and the logarithmic power spectra of the flares and the quiet state star deflection. The logarithmic power spectrum will enable us to separate the random part of the spectrum from the non random. (2) We isolate by filtering the random part of the spectrum and we estimate the standard deviation of the random noise σ_{BW} . (3) We identify the potential frequencies of star brightness oscillations from the power spectrum. To do this we use the Ho hypothesis test (see [1]) in which $\sigma = \sigma_{BW}$. (4) We filter out the identified frequencies from the star deflection. (5) We estimate the confidence level of those frequencies identifications comparing the amplitudes of the oscillations with the respective σ_{BW} .

4 Results

DFT-analysis and the use of the Brownian Walk noise notion enable us to estimate the proper random noise and to de-noise by filtering the observed light-curves. Thus we were able to detect possible weak transient optical oscillations and to isolate them by proper band filtering with a confidence level lower than 30% (Probability of proper identification higher than 70%). The results for all the program stars indicate that: (1) Transient high frequency oscillations occur during the quiet-star phase. (2) The Observed frequencies range between 0.0083Hz (period 2min) and 0.3 Hz (period 3s) not rigorously bounded. The observed oscillations are of the same frequencies range of the sunquakes which are high-degree, high-frequency p modes waves (standing acoustic waves [11])

References

- [1] Contadakis, M. E., Avgoloupis, S. J., Seiradakis J. H., et al., 2004, *Astron. Nachr.*, 325, No. 5, pp. 428-432.
- [2] Contadakis, M.E., Avgoloupis, S. J., Seiradakis J. H., 2006, Further investigation of the high frequency optical oscillations during the flare phase of the red dwarf EV Lac in *Recent Advances in Astronomy and Astrophysics*, Edited by N. Solomos , Proceedings of the 7nth International Conference of HELLAS, AIP press, 324
- [3] Contadakis, M.E., Avgoloupis, S.J. and Seiradakis, J.H., 2010, Transient high frequency optical oscillations of the red dwarf AD Leonis. In *Advances in Hellenic Astronomy during the IY A09*, eds. K.Tsinganos, D.Hatzidimitriou, T.Matsakos, ASPCS Vol. 424, pp189-191, 189
- [4] Contadakis, M.E., Avgoloupis, S.J., and Seiradakis, J.H., 2011b, Transient high frequency optical oscillations of the flare star AD Leonis, in *The Apple of the Knowledge*, Eds M.E.Contadakis, C.Kaltsikis, S.Spatalas, K.Tokmakidis and I.N.Tziavos, Ziti Press Thessaloniki, Greece, 459
- [5] Contadakis, M.E. Avgoloupis, S.J., Seiradakis, J.H.: 2011b, JENAM-2011, 4-8 July 2011, Saint Petersburg, Russia
- [6] Contadakis, M.E., Avgoloupis, S.J., Seiradakis, J.H.:2011c, JENAM-2011, 4-8 July 2011, Saint Petersburg, Russia
- [7] Contadakis, M.E. , Avgoloupis, S.J., Seiradakis, J.H.: 2011d, 10th Hellenic Astronomical Conference 5-8 September 2011, Ioannina, Greece
- [8] Contadakis , M.E. Avgoloupis, S.J., Seiradakis, J.H.:2011e, 10th Hellenic Astronomical Conference 5-8 September 2011, Ioannina, Greece
- [9] Contadakis, M.E. Avgoloupis, S.J., Seiradakis, J.H.: 2012, *AN* 333, No. 7, 583 - 593 (2012) / DOI 10.1002/asna.201111690
- [10] Contadakis, M.E. , Avgoloupis, S.J., Seiradakis, J.H.: 2013, *AApTr* vol. 28,1, 9-14
- [11] Kosovichev, A.G., 2013, *Proceedings IAU Symposium on Precision Astroseismology*, Vol.301, pp 349-352
- [12] Zhilyaev, B.E., Romanyuk, Ya.O., Verlyuk, I.A., et al. , 2007, *A & A*, 465, 235